

20832  
5746-7438

# WORLD HEALTH STATISTICS 2007



World Health  
Organization

**WHO Library Cataloguing-in-Publication Data**

World health statistics 2007.

1. Health status indicators. 2. World health. 3. Health services - statistics. 4. Mortality.  
5. Life expectancy. 6. Demography. 7. Statistics. I. World Health Organization.

ISBN 978 92 4 156340 6  
ISBN 978 92 4 068211 5 (electronic version)

(NLM classification: WA 900.1)

**© World Health Organization 2007**

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; e-mail: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

This publication was produced by the Department of Measurement and Health Information Systems of the Information, Evidence and Research Cluster, under the overall direction of Ties Boerma and Kenji Shibuya, in collaboration with WHO technical programmes and regional offices, and assisted by Zoe Brillantes, Maria Guraib, Mie Inoue, Yohannes Kinfu and Doris Ma Fat.

Valuable inputs to the statistical highlights in Part 1 were received from Monika Bloessner, Ties Boerma, Somnath Chatterji, Mercedes de Onis, Christopher Dye, Christopher Fitzpatrick, Charu Garg, Mehran Hosseini, Ahmadreza Hosseinpoor, Mie Inoue, Yohannes Kinfu, Doris Ma Fat, Colin Mathers, Ritu Sadana, Kenji Shibuya, Tessa Tan-Torres and Catherine Watt. Maps were produced by the Public Health Mapping and Geographic Information Systems team, Communicable Disease and Surveillance.

Contributors to the statistical tables in Part 2 were: Michel Beusenbergh, Monika Bloessner, Cynthia Boschi Pinto, Claire Chauvin, Mercedes de Onis, Christopher Dye, Christopher Fitzpatrick, Marta Gacic Dobo, Charu Garg, Chika Hayashi, Mehran Hosseini, Ahmadreza Hosseinpoor, Chandika Indikadahena, Mie Inoue, Yohannes Kinfu, Teena Kunjumen, Doris Ma Fat, Colin Mathers, Chizuru Nishida, Vladimir Pozniak, Eva Rehfuss, Dag Rekve, Leanne Riley, Lale Say, Kenji Shibuya, Jonathan Siekmann, Jacqueline Sims, Yves Souteyrand, Tessa Tan-Torres, Jeanette Vega, Catherine Watt, and many staff in WHO country offices, governmental departments and agencies and international institutions. Additional help and advice were kindly provided by regional offices and members of their staff, including Yok-Ching Chong, Anton Fric, Remigijus Prochorskas, Saher Shuqaidef, William Soumbey-Alley and Fernando Zacarias.

The publication was edited by Miriam Pinchuk. Editorial and production support was provided by the Department of Knowledge Management and Sharing, including Caroline Allsopp, Ian Coltart, Laragh Golligly, Maryvonne Grisetti, Sophie Guetaneh Aguetant, Hooman Momen, and Catherine Roch. The web site version and other electronic media were provided by the Digital Publishing Solution, Ltd. Proofreading was by Melanie Lauckner. We also thank Susan Piccolo and Petra Schuster for their administrative support.

Printed in France



# Table of Contents

<b>Introduction</b>	<b>7</b>
<b>Part 1. Ten statistical highlights in global public health</b>	<b>9</b>
1. Monitoring progress: appropriate use of health statistics	10
2. People living with HIV: better data, better estimates	11
3. Future health: projected deaths for selected causes to 2030	12
4. Child undernutrition: where are we now?	13
5. Levels and causes of death: filling data gaps	14
6. Tobacco use and poverty: high prevalence among the world's poorest	15
7. Mental illness: depression worsens the health of people with chronic illness	16
8. Inequalities in health: understanding their determinants	17
9. Tuberculosis control: towards goals and targets	18
10. Health expenditure: meeting needs?	19
References	20
<b>Part 2. World health statistics</b>	<b>21</b>
<b>Health status: mortality</b>	<b>22</b>
Life expectancy at birth (years)	
Healthy life expectancy (HALE) at birth (years)	
Probability of dying aged 15–60 years per 1 000 population (adult mortality rate)	
Probability of dying aged < 5 years per 1 000 live births (under-5 mortality rate)	
Infant mortality rate (per 1 000 live births)	
Neonatal mortality rate (per 1 000 live births)	
Maternal mortality ratio (per 100 000 live births)	
Deaths due to HIV/AIDS (per 100 000 population per year)	
Deaths due to tuberculosis among HIV-negative people (per 100 000 population per year)	
Deaths due to tuberculosis among HIV-positive people (per 100 000 population per year)	
Age-standardized mortality rate by cause (per 100 000 population)	
Distribution of years of life lost by broader causes (%)	
Distribution of causes of death among children aged < 5 years (%)	
<b>Health status: morbidity</b>	<b>32</b>
HIV prevalence among adults aged ≥ 15 years (per 100 000 population)	
Prevalence of tuberculosis (per 100 000 population)	
Incidence of tuberculosis (per 100 000 population per year)	
Number of confirmed cases of poliomyelitis	



# Table of Contents

## **Health service coverage** **36**

---

- Immunization coverage among 1-year-olds with one dose of measles (%)
- Immunization coverage among 1-year-olds with three doses of diphtheria, tetanus toxoid and pertussis (DTP3) (%)
- Immunization coverage among 1-year-olds with three doses of Hepatitis B (HepB3) (%)
- Antenatal care coverage (%)
- Births attended by skilled health personnel (%)
- Contraceptive prevalence rate (%)
- Children aged < 5 years sleeping under insecticide-treated bednets (%)
- Antiretroviral therapy coverage among people with advanced HIV infections (%)
- HIV-infected pregnant women who received antiretrovirals for PMTCT (%)
- Tuberculosis detection rate under DOTS (%)
- Tuberculosis treatment success under DOTS (%)
- Children aged < 5 years with ARI symptoms taken to facility (%)
- Children aged < 5 years with diarrhoea receiving ORT (%)
- Children aged < 5 years with fever who received treatment with any antimalarial (%)
- Children 6–59 months who received vitamin A supplementation (%)
- Births by Caesarean section (%)

## **Risk factors** **46**

---

- Children aged < 5 years stunted for age (%)
- Children aged < 5 years underweight for age (%)
- Children aged < 5 years overweight for age (%)
- Low-birthweight newborns (%)
- Adults aged ≥ 15 years who are obese (%)
- Access to improved drinking water sources (%)
- Access to improved sanitation (%)
- Population using solid fuels (%)
- Prevalence of current tobacco use in adolescents (13–15 years) (%)
- Prevalence of current tobacco use among adults (≥ 15 years) (%)
- Per capita recorded alcohol consumption (litres of pure alcohol) among adults (≥ 15 years)
- Prevalence of condom use by young people (15–24 years) at higher risk sex (%)

## **Health systems** **56**

---

- Human resources for health 56
  - Physicians; Nurses; Midwives; Dentists; Pharmacists; Public and environmental health workers; Community health workers; Laboratory health workers; Other health workers; Health management and support workers



# Table of Contents

Health expenditure ratios	64
Total expenditure on health as % of gross domestic product	
General government expenditure on health as % of total expenditure on health	
Private expenditure on health as % of total expenditure on health	
General government expenditure on health as % of total government expenditure	
External resources for health as % of total expenditure on health	
Social security expenditure on health as % of general government expenditure on health	
Out-of-pocket expenditure as % of private expenditure on health	
Private prepaid plans as % of private expenditure on health	
Health expenditure aggregates	
Per capita total expenditure on health at average exchange rate (US\$)	65
Per capita total expenditure on health at international dollar rate	
Per capita government expenditure on health at average exchange rate (US\$)	
Per capita government expenditure on health at international dollar rate	
Coverage of vital registration of deaths (%)	65
Hospital beds (per 10 000 population)	65

## **Inequities in health** **74**

Probability of dying aged < 5 years per 1 000 live births (under-5 mortality rate) by place of residence; by wealth quintile; by educational level of mother	
Children aged < 5 years stunted for age (%) by place of residence; by wealth quintile; by educational level of mother	
Births attended by skilled health personnel (%) by place of residence; by wealth quintile; by educational level of mother	
Measles immunization coverage among 1-year-olds by place of residence; by wealth quintile; by educational level of mother	

## **Demographic and socioeconomic statistics** **78**

Population (thousands)	
Annual population growth rate (%)	
Population in urban areas (%)	
Total fertility rate (per woman)	
Adolescent fertility rate (%)	
Adult literacy rate (%)	
Net primary school enrolment ratio (%)	
Gross national income per capita (international\$)	
Population living below the poverty line (% living on < US\$1 per day)	





# Introduction

*World health statistics 2007* presents the most recent health statistics for WHO's 193 Member States. This third edition includes a section with 10 highlights of global health statistics for the past year as well as an expanded set of 50 health statistics.

*World health statistics 2007* has been collated from publications and databases produced by WHO's technical programmes and regional offices. The core set of indicators was selected on the basis of their relevance to global health, the availability and quality of the data, and the accuracy and comparability of estimates. The statistics for the indicators are derived from an interactive process of data collection, compilation, quality assessment and estimation occurring among WHO's technical programmes and its Member States. During this process, WHO strives to maximize the accessibility, accuracy, comparability and transparency of health statistics.<sup>1</sup>

In addition to national statistics, this publication presents statistics on the distribution of selected health outcomes and interventions within countries, disaggregated by gender, age, urban versus rural setting, wealth, and educational level. Such statistics are primarily derived from analyses of household surveys and are available only for a limited number of countries. We envisage that the number of countries reporting disaggregated data will increase during the next few years.

The core indicators do not aim to capture all relevant aspects of health but to provide a comprehensive summary of the current status of a population's health and the health system at country level. These indicators include: mortality outcomes, morbidity outcomes, risk factors, coverage of selected health interventions, health systems, inequalities in health, and demographic and socioeconomic statistics.

All statistics have been cleared as WHO's official figures in consultation with Member States unless otherwise stated. WHO's estimates use data from publicly accessible databases, peer-reviewed methods of estimation, and consultation with experts around the world. The estimates published here should, however, still be regarded as best estimates made by WHO rather than the official view of Member States.

As the demand for timely, reliable and comparable information on key health statistics continues to increase, users need to be well informed about the definitions used and the quality and limitations of health statistics. More detailed information, including a compendium of statistics and an online version of this publication, is available from WHO's Statistical Information System (<http://www.who.int/statistics>). The web site also includes information on how each statistic is derived.

The online version of *World health statistics 2007* will be updated regularly, and it includes the most recent estimates and time-series of relevant health statistics. The online version also provides, whenever possible, metadata describing the sources of data, estimation methods and quality of estimates. It is hoped that careful scrutiny and use of the statistics presented in this report will lead to progressively better measurement of relevant indicators of population health and health systems.

---

1. To meet these objectives, WHO has initiated the organization-wide Programme on Health Statistics. For more information, see <http://www.who.int/healthinfo/statistics/programme/en/index.html>.








## Part 1

---

# Ten statistical highlights in global public health



# 1. Monitoring progress: appropriate use of health statistics



The ability to monitor progress towards the Millennium Development Goals (MDGs) depends primarily on data availability. There is a stark contrast between the data available about the under-five mortality rate, the indicator for MDG 4, and the maternal mortality ratio, against which MDG 5 is monitored.


Under-five mortality rates are derived from vital registration systems, censuses and household surveys.<sup>1</sup> In most countries, there are data points available over time, and these are analysed to obtain the best current estimate. Uncertainty occurs when there is a need to project estimates forward to the current year since the most recent data generally refer to a few years earlier. Measuring the maternal mortality ratio has been a greater challenge because, compared with deaths among children, maternal deaths are rare events. In countries without a complete death registration system and medical certification, large-scale household surveys or censuses using verbal autopsy techniques provide estimates of the ratio, since facility-based statistics are inherently biased. Even then, much uncertainty remains. As a consequence, the global estimate of the maternal mortality ratio is published only once every five years, and in 2000, 40% of countries' estimates were based on figures predicted by regression.<sup>4</sup> The ability to reliably assess trends in maternal mortality is limited.

For monitoring, it is important to distinguish between corrected and predicted statistics.<sup>5,6</sup> Corrected statistics use adjustments made for known biases and, if needed, are based on a systematic reconciliation of data from multiple sources using established, transparent methods. Predicted statistics use a set of assumptions about the association between other factors and the quantity of interest, such as maternal mortality, to fill gaps in the data over time (projecting into the present or future) or space (from one population with data to another with limited or no data). Predicted statistics are not suitable for monitoring progress. Unfortunately, the MDG monitoring process relies heavily on predicted statistics.<sup>5</sup> This mismatch was created partly by the demand for more timely statistics and partly by the lack of data and good measurement strategies for certain statistics. It is crucial for the international community to invest in data collection and use indicators that are valid, reliable and comparable; the international community must also have well-defined measurement strategies for monitoring progress and evaluating health programmes.<sup>7</sup>



## 2. People living with HIV: better data, better estimates

**Number of people living with HIV: comparing past and current estimates<sup>8-14</sup>**



The exact number of people living with HIV is unknown despite the fact that HIV infection can easily be diagnosed by a widely used antibody test. Achieving 100% certainty about the number of people living with HIV globally would require testing every person in the world for HIV every year. Nonetheless, we can estimate the number by using data from different sources, such as surveillance of pregnant women attending antenatal clinics, household surveys with HIV testing and sentinel surveillance among populations at higher risk of HIV infection.

UNAIDS and WHO, in close consultation with countries, employ a standardized method for obtaining estimates of HIV prevalence among men and women. An increasing number of countries have adopted these methods to develop their own national estimates. But an estimate is only as good as the data. As more complete data become available, past estimates may need to be adjusted. This is the case for the AIDS epidemic. The bars in the figure estimate the number of people infected with HIV at the time of publication of each annual *AIDS epidemic update* since 2000.<sup>8-14</sup> The line shows the best estimates for each year that were made in 2006 in the most recent update: this reveals not only that the size of the epidemic had been overestimated previously but also that it is still growing. The ranges around the estimates reflect the degree of uncertainty about global HIV estimates.


Improvements in recent estimates are the result of revisions made using better data. These revisions used data from national population-based surveys and benefited from improvements in the quality and coverage of sentinel surveillance systems in many countries.

The latest estimates cannot be compared directly with estimates published previously. It would be incorrect to derive a trend by comparing the bars. The 2006 estimates for this year and past years (indicated by the line) are more accurate than those produced in previous years since they are based on improved methods and used more data than earlier estimates. The need to exercise caution is not unusual when comparing global estimates of disease over time.



### 3. Future health: projected deaths for selected causes to 2030

Projected global deaths for selected causes of death, 2002–2030<sup>15</sup>



Predicted statistics have an important and useful role in helping to inform planning and strategic decision-making, and in prioritizing research and development issues. According to projections carried out by WHO and published in early 2006,<sup>15</sup> the world will experience a substantial shift in the distribution of deaths from younger age groups to older age groups, and from communicable diseases to noncommunicable diseases during the next 25 years. Large declines in mortality are projected to occur between 2002 and 2030 for all of the principal communicable, maternal, perinatal and nutritional causes, with the exception of HIV/AIDS. Global deaths from HIV/AIDS are projected to rise from 2.8 million in 2002 to 6.5 million in 2030 under a baseline scenario that assumes antiretroviral drug coverage reaches 80% by 2012.


Although age-specific death rates for most noncommunicable diseases are projected to decline, the ageing of the global population will result in significant increases in the total number of deaths caused by most non-communicable diseases over the next 30 years. Overall, noncommunicable conditions will account for almost 70% of all deaths in 2030 under the baseline scenario. The projected 40% increase in global deaths resulting from injury between 2002 and 2030 is predominantly due to the increasing number of deaths from road traffic accidents.

The four leading causes of death globally in 2030 are projected to be ischaemic heart disease, cerebrovascular disease (stroke), HIV/AIDS and chronic obstructive pulmonary disease. The total number of tobacco-attributable deaths is projected to rise from 5.4 million in 2005 to 6.4 million in 2015 and to 8.3 million in 2030. Tobacco is projected to kill 50% more people in 2015 than HIV/AIDS and to be responsible for 10% of all deaths.



## 4. Child undernutrition: where are we now?

### Geographical pattern of stunting in children under 5 years of age<sup>16</sup>



The release of WHO's new Child Growth Standards (<http://www.who.int/childgrowth/en>) has an impact on estimates of undernutrition among children. Global, regional and country estimates have been recalculated using the new standards, which include data from 388 national surveys in 139 countries.<sup>16</sup>


In 2005, in all developing countries 32% of children under 5 years of age (178 million children) were estimated to be stunted (that is, their height fell –2 standard deviations below the median height-for-age of the reference population). In that year, more than 40% of stunting was found in the WHO regions of Africa and South-East Asia, around 25% in the Eastern Mediterranean Region and 10–15% in the regions of the Americas and the Western Pacific. Of the 39 countries with a prevalence of stunting of 40% and higher, 22 are in the African Region, 7 in South-East Asia, 4 in the Eastern Mediterranean, 4 in the Western Pacific, and 1 each in Europe and in the Americas. Of the 35 countries with a stunting prevalence lower than 20%, 13 are in the Region of the Americas, 11 in Europe, 6 in the Eastern Mediterranean, 3 in the Western Pacific and 2 in South-East Asia.

Wasting (defined as being –2 standard deviations below the median of weight-for-height) is a sign of acute malnutrition and is a strong predictor of mortality among children. The global estimate of wasting occurring among children under 5 years of age based on WHO's new standards is 10% (or 55 million). The highest number of affected children – 29 million – is estimated to live in south-central Asia. The same regional pattern is found for severe wasting (defined as being –3 standard deviations below the median), with an estimated total prevalence of 4% – or 19 million – children affected. Many of these children are likely to die before reaching the age of 5 years. In general, compared with estimates based on the previous international reference, stunting rates are higher for all age groups when the new WHO standards are used. Additionally, the prevalences of wasting and severe wasting are higher during the first half of infancy with the new WHO standards; and thereafter severe wasting rates continue to be 1.5 to 2.5 times higher than those of the previous reference.



## 5. Levels and causes of death: filling data gaps

### Quality of cause-of-death information from national civil registration systems, based on latest data received from WHO Member States, circa 2003<sup>17,19</sup>



**Note:** The criteria used to assess the quality of cause-of-death information are valid for data from national civil registration systems. Therefore, they do not apply to China and India since they report data from sample registration systems, which cover < 10% of their populations.

Accurate and timely data on deaths and causes of death with medical certification are essential. WHO collects information on causes of death from its Member States annually. However, for more than a fourth of the world's population – largely located in Africa, South-East Asia and the Middle East – there are no recent data available to WHO, and these are the areas where much of the burden of disease falls. Altogether, 115 Member States have some form of death registration known to WHO; this includes China and India, which also have sample vital registration systems.

There are delays in compiling, analysing and reporting these statistics: by 2007 WHO had received reports for 2004 or 2005 for 64 (56%) countries. An assessment of the quality of cause-of-death information by WHO suggested that ideal systems operate in only 29 of 115 countries that report such statistics to WHO; these systems represent less than 13% of the world's population.<sup>17</sup> In the remaining countries mortality statistics suffer from one or more of the following problems: incomplete registration of births and deaths, lay reporting of the cause of death, poor coverage and incorrect reporting of ages.


The ultimate goals should be to establish complete vital registration with medical certification of deaths in all countries. National governments, with the support of international organizations, need to continue to make efforts to improve the coverage and quality of vital registration systems.

At the same time, complementary approaches to complete vital registration are needed to respond to the demand for timely information and to assess the performance of the systems themselves. WHO, in collaboration with its partners, is stepping up efforts to improve the quality of data that underlies its overall estimates of mortality by age, sex and cause.<sup>18</sup> Such efforts include making better use of household surveys and censuses, implementing standardized verbal autopsy instruments, and using data from partial vital registration and sources other than vital registration.



## 6. Tobacco use and poverty: high prevalence among the world's poorest

**Daily tobacco smoking among adults aged 18 years and older,  
by income quintile and WHO region<sup>20</sup>**



SEAR, South-East Asia; AFR, African; EMR, Eastern Mediterranean; WPR, Western Pacific; AMR, Americas; EUR, European.

Health inequalities refer to differences in health status or in the distribution of health determinants between different populations. The burden of disease attributable to tobacco use weighs increasingly heavily on populations in developing economies. According to the latest estimates, more than 80% of the 8.3 million deaths attributed to tobacco and projected to the year 2030 will occur in low-income and middle-income countries.<sup>15</sup>

Data on the prevalence of smoking among adults in developing countries are limited. WHO's World Health Survey provides a valuable insight into the comparative prevalence among adults aged 18 and older.<sup>20</sup> The results of the 2003–2004 survey indicate that daily tobacco smoking is most prevalent among the lowest-income households in developing economies – that is, among the poorest of the poor. Indeed, prevalence is highest among the poor in all WHO regions except the European Region. The difference in prevalence between the poor and the (relatively) rich is greatest among the group of South-East Asian countries surveyed, where average per capita income is lowest.

The combination of a higher prevalence of tobacco use and more limited access to health resources results in severe health inequalities, and is likely to perpetuate the vicious circle of illness and poverty. Inequalities between and within countries in terms of the risk of infectious diseases now have been extended to inequalities in risk factors for noncommunicable diseases; this has implications for health systems at all levels.


\* Surveyed countries in each region include: African Region (AFR): Burkina Faso, Chad, Comoros, Congo, Côte d'Ivoire, Ethiopia, Ghana, Kenya, Malawi, Mali, Mauritania, Mauritius, Namibia, Senegal, South Africa, Swaziland, Zambia, Zimbabwe; Region of the Americas (AMR): Brazil, Dominican Republic, Ecuador, Guatemala, Mexico, Paraguay, Uruguay; Eastern Mediterranean Region (EMR): Morocco, Pakistan, Tunisia, United Arab Emirates; European Region (EUR): Bosnia and Herzegovina, Croatia, Czech Republic, Estonia, Georgia, Hungary, Kazakhstan, Latvia, Russian Federation, Slovakia, Slovenia, Spain, Ukraine; South-East Asia Region (SEAR): Bangladesh, India, Sri Lanka, Myanmar, Nepal; Western Pacific Region (WPR): China, Laos, Malaysia, Philippines, Viet Nam.





## 7. Mental illness: depression worsens the health of people with chronic illness

Mean health score by disease status, World Health Survey 2003<sup>20</sup>



Depression is an important global public health problem due to both its relatively high lifetime prevalence and the significant disability that it causes. In 2002, depression accounted for 4.5% of the worldwide total burden of disease (in terms of disability-adjusted life years). It is also responsible for the greatest proportion of burden attributable to non-fatal health outcomes, accounting for almost 12% of total years lived with disability worldwide.<sup>21</sup> Without treatment, depression has the tendency to assume a chronic course, to recur, and to be associated with increasing disability over time.

WHO's World Health Survey collected data on health and health-related outcomes and their determinants in samples of adults aged 18 years and older.<sup>20</sup> The prevalence of depression was estimated using criteria in the *International statistical classification of diseases and related health problems, tenth revision* (ICD-10). The prevalences of four chronic physical diseases – angina, arthritis, asthma and diabetes – were also estimated. The figure shows the mean health score – where 0 is the worst level of health and 100 is the best level of health – for each disease with and without accompanying depression. Individuals without depression and without other conditions had a mean health score of 90. Respondents with only one of the chronic diseases had mean health scores of around 80. Respondents with depression but without chronic disease had the lowest mean health score (73). Respondents with depression and another chronic condition had much lower mean health scores when compared with respondents who had only a chronic condition. These patterns were consistent after adjusting for sociodemographic variables.


This analysis does not tell us whether people are more depressed because they have a coexisting chronic condition. The timely diagnosis and treatment of depressive disorders are essential irrespective of causality. In many primary care settings when patients present with multiple disorders that include depression, the depression often remains undiagnosed, and even if it is diagnosed, treatment usually focuses on the other chronic diseases. Depression can be treated in primary care or community settings using locally available and cost-effective interventions.






## 8. Inequalities in health: understanding their determinants

**Stunting among children under 5 years of age, by household wealth quintile, Mozambique, 1999–2003<sup>23</sup>**



**What contributes to inequity in childhood stunting?<sup>23</sup>**



**Note:** Household wealth index constructed using durable goods, type of materials used in housing floor and number of rooms divided by the number of household members. Wealth quintile 1 indicates the poorest and wealth quintile 5, the least poor.

Measuring socioeconomic inequalities in a population's health is important because national averages often mask differences within and across subgroups. For policy purposes it is especially relevant to understand why unfair and avoidable inequalities (or inequities) exist and what actions may be taken to improve equity. Decomposition analysis is one approach used to quantify the contribution made by different factors to inequities in health; it takes into account the socioeconomic distribution of determinants of health and health indicators.<sup>22</sup> Such analysis can serve as one input to aid in the development of evidence-based policies, relevant to a particular context or country, to reduce inequities.

For example, decomposition analysis using data from the 2003 Demographic and Health Survey in Mozambique shows that the four biggest contributors to poor growth in children (defined as height-for-age falling 2 standard deviations below the median of the reference population) stratified by household wealth are: source of drinking water (19%), household wealth itself (17%), geographical differences (16%) and mother's occupation (13%).<sup>23</sup> An additional 10 factors identified in the survey together contribute 35%. Using this technique to uncover inequities reveals that strategies to address contributing factors are likely to require collaborative and intersectoral actions that are not limited to health authorities or the health system.


Describing health inequities and understanding their determinants require process and outcome data that can be disaggregated by different socioeconomic or demographic characteristics, as well as the ability to link data from different sectors in a country. WHO is contributing to these efforts by setting norms and standards, and providing technical assistance to Member States.




## 9. Tuberculosis control: towards goals and targets

### 26 million TB patients treated but global targets narrowly missed in 2005<sup>24</sup>

**Global case detection 60% in 2005**  
Target reached in Western Pacific Region



**Global treatment success 84% in 2004–2005**  
Target reached in South-East Asia and Western Pacific regions



There were an estimated 8.8 million new tuberculosis (TB) cases in 2005, including 7.4 million in Asia and sub-Saharan Africa. A total of 1.6 million people died of TB, including 195 000 patients infected with HIV. Using surveillance data, *Global tuberculosis control: surveillance, planning, financing* draws four main conclusions about TB control programmes.<sup>24</sup>


First, although more than 26 million TB patients have been treated under WHO's DOTS strategy, the world's TB control programmes narrowly missed their 2005 targets for case detection (reaching 60% compared to the target of 70%) and cure (84% compared to the target of 85%). However, both targets were met in WHO's Western Pacific Region and in 26 countries including China, the Philippines and Viet Nam. Second, while the total number of patients diagnosed and treated in 2005 using DOTS approached the target, the number of patients known to be HIV positive or carrying multidrug-resistant TB (MDR-TB) were far fewer than anticipated by *The Global Plan to Stop TB 2006–2015*.<sup>25</sup> Therefore, major efforts are needed to step up collaborative activities between TB and HIV programmes and to manage MDR-TB and extensively drug-resistant TB. Third, the global TB epidemic appears to be on the threshold of decline. The incidence rate is now stable or falling in all WHO regions, including Africa and Europe.

These findings, if robust, mean that MDG Target 8 ("Have halted by 2015 and begun to reverse the incidence of malaria and other major diseases [including TB]") will be met before 2015. However, the total number of new cases was still rising slowly in 2005 in WHO's African, Eastern Mediterranean and South-East Asia regions. For reasons that are not fully understood, in Asian countries that report high rates of case detection and treatment success, the incidence has not been reduced as quickly as expected. This is linked to the fourth conclusion: the global burden of TB is not falling fast enough to satisfy the more demanding targets set by the Stop TB Partnership. At the current pace, 1990's prevalence and mortality rates will not be halved worldwide by 2015.



## 10. Health expenditure: meeting needs?

### Percentage distribution of population, disability-adjusted life years (DALYs) and total health expenditure by WHO region and membership of Organisation for Economic Co-operation and Development (OECD), 2004<sup>26</sup>



AFR, African; AMR, Americas; EMR, Eastern Mediterranean; EUR, European; SEAR, South-East Asia; WPR, Western Pacific.

**Note:** Totals for the following regions calculated after subtracting the 30 OECD members: Americas, European and Western Pacific. DALYs are from 2002.

In 2004, the world spent a total of US\$ 4.1 trillion on health, which is equivalent to 4.9 trillion international dollars. (International dollars are used to account for the purchasing power of different national currencies.) The geographical distribution of financial resources for health is uneven.<sup>26</sup> There is a 20/90 syndrome in which 30 member countries of the Organisation for Economic Co-operation and Development (OECD) make up less than 20% of the world's population but spend 90% of the world's resources on health.

OECD countries spend a larger share of their gross domestic product on health, spending on average more than 11%, compared with 4.7% for countries in WHO's African and South-East Asia regions. This translates to per capita spending of about 3080 international dollars (US\$ 3170) in OECD countries compared with 102 international dollars (US\$ 36) in countries in the African and South-East Asia regions, which are much poorer. Linking this spending to epidemiology, the figure shows that although poorer WHO regions, such as Africa and South-East Asia, account for the largest share of the global burden of disease (more than 50% of global disability-adjusted life years lost) and 37% of the world's population, they spend about 2% of global resources on health. The Western Pacific Region, excluding Australia, Japan, New Zealand and the Republic of Korea, accounts for 24% of the world's population (which is dominated by China), about 18% of the global burden of disease but only 2% of the world's health resources. The Region of the Americas and the European Region, excluding the OECD countries, account for about 12% of the world's population, 11% of the global burden of disease and spend slightly less than 5% of health resources.

Richer countries with smaller populations and lower disease burdens use more health resources than poorer countries with larger populations and higher disease burdens. This highlights the absolute need for additional resources for many poor countries and raises questions about the efficiency of spending on health in richer countries.



## References

1. Hill K et al. Trends in child mortality in the developing world: 1990 to 1996. New York, UNICEF, 1998.
2. United Nations Children's Fund. *State of the world's children 2007*. New York, United Nations Children's Fund, 2006.
3. ORC Macro. *Demographic and health survey: Malawi 2007*. Calverton, MD, ORC Macro, 2007.
4. *Maternal mortality in 2000: estimates developed by WHO, UNICEF and UNFPA*. Geneva, World Health Organization, 2004.
5. Murray CJ. Towards good practice for health statistics: lessons from the Millennium Development Goal health indicators. *Lancet*, 2007, 369:862–873.
6. *Advisory Committee on Health Monitoring and Statistics: meeting report*. Geneva, World Health Organization, 2006 (<http://www.who.int/healthinfo/statistics/healthinfoachmsreport20061214-15.pdf>, accessed 4 April 2007).
7. Boerma JT, Stansfield SK. Health statistics now: are we making the right investments? *Lancet*, 2007, 369:779–786.
8. *AIDS epidemic update: December 2000*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2000 (WHO/CDS/CSR/EDC/2000.9).
9. *AIDS epidemic update: December 2001*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2001 (WHO/CDS/CSR/NCS/2001.2).
10. *AIDS epidemic update: December 2002*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2002 (UNAIDS/02.58E).
11. *AIDS epidemic update: December 2003*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2003 (UNAIDS/03.39E).
12. *AIDS epidemic update: December 2004*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2004 (UNAIDS/04.16E).
13. *AIDS epidemic update: December 2005*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2005 (UNAIDS/05.19E).
14. *AIDS epidemic update: December 2006*. Geneva, Joint United Nations Programme on HIV/AIDS, World Health Organization, 2006 (UNAIDS/06.29E).
15. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Medicine* [online journal], 2006, 3(11):e442 (<http://medicine.plosjournals.org/perlserv/?request=get-document&doi=10.1371/journal.pmed.0030442>, accessed 4 April 2007).
16. Global database on child growth and malnutrition [online database]. Geneva, World Health Organization, 2007 (<http://www.who.int/nutgrowthdb/database/en>, accessed 4 April 2007).
17. Mathers CD et al. Counting the dead and what they died from: an assessment of the global status of cause of death data. *Bulletin of the World Health Organization*, 2005, 83:171–177.
18. Shibuya K. Counting the dead is essential for health. *Bulletin of the World Health Organization*, 2006, 84:170–171.
19. *WHO mortality database: tables* [online database]. Geneva, World Health Organization, 2007 (<http://www.who.int/healthinfo/mortables/en/index.html>, accessed 4 April 2007).
20. *WHO survey data centre: World Health Survey*. Geneva, World Health Organization, 2007 (<http://surveydata.who.int/>, accessed 4 April 2007).
21. *Revised global burden of disease (GBD) 2002 estimates*. Geneva, World Health Organization, 2005 (<http://www.who.int/healthinfo/bodgbd2002revised/en/index.html>, accessed 4 April 2007).
22. Hosseinpoor AR et al. Decomposing socioeconomic inequality in infant mortality in Iran. *International Journal of Epidemiology*, 2006, 35:1211–1219.
23. *A WHO report on inequities in maternal and child health in Mozambique*. Geneva, World Health Organization, 2007.
24. *Global tuberculosis control: surveillance, planning, financing. WHO report 2007*. Geneva, World Health Organization, 2007 (WHO/HTM/TB/2007.376).
25. *The Global Plan to Stop TB 2006–2015*. Geneva, Stop TB Partnership, World Health Organization, 2006 (WHO/HTM/STB/2006.35).
26. *National health accounts*. Geneva, World Health Organization, 2007 (<http://www.who.int/nha>, accessed 4 April 2007).